



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 31/505, 31/44, 31/35, 31/225	A1	(11) International Publication Number: WO 98/11896 (43) International Publication Date: 26 March 1998 (26.03.98)
(21) International Application Number: PCT/US97/16388 (22) International Filing Date: 15 September 1997 (15.09.97) (30) Priority Data: 60/026,581 18 September 1996 (18.09.96) US 9621970.4 22 October 1996 (22.10.96) GB (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): GOULD, Robert, J. [CA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). NICHTBERGER, Steven, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). RHYMER, Patricia, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). OLOFSSON, Lars [SE/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: COMBINATION THERAPY FOR REDUCING THE RISKS ASSOCIATED WITH CARDIOVASCULAR DISEASE (57) Abstract The instant invention involves a combination therapy and pharmaceutical compositions comprised of a therapeutically effective amount of a cholesterol reducing agent such as an HMG-CoA reductase inhibitor in combination with a platelet aggregation inhibitor which is useful for inhibiting platelet aggregation, for inhibiting the formation of thrombotic occlusions, and for treating, preventing and reducing the risk of occurrence of cardiovascular and cerebrovascular events and related vaso-occlusive disorders.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

TITLE OF THE INVENTION

COMBINATION THERAPY FOR REDUCING THE RISKS
ASSOCIATED WITH CARDIOVASCULAR DISEASE.

5 FIELD OF THE INVENTION

The instant invention involves a combination therapy comprising the administration of a cholesterol reducing agent such as a 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitor (or HMG-CoA RI) and a platelet aggregation inhibitor for
10 treating, preventing and reducing the risk of developing cardiovascular and cerebrovascular events and disorders in a mammal.

BACKGROUND OF THE INVENTION

Platelet activation and aggregation are involved in unstable
15 angina and acute myocardial infarction, in reocclusion following thrombolytic therapy and angioplasty, in transient ischemic attacks and in a variety of other vaso-occlusive disorders. When a blood vessel is damaged either by acute intervention such as angioplasty, or more chronically by the pathophysiological processes of atherosclerosis,
20 platelets are activated to adhere to the disrupted surface and to each other. This activation, adherence and aggregation may lead to occlusive thrombus formation in the lumen of the blood vessel.

Antiplatelet therapy has been used in a wide variety of cardiovascular disease states and in conjunction with interventional
25 therapy such as coronary artery or peripheral bypass grafting, cardiac valve replacement, and percutaneous transluminal coronary angioplasty (PTCA). Available drugs, such as aspirin and ticlopidine (TICLID®), have shown efficacy in syndromes involving vascular occlusion, presumably due to sustained inhibition of platelet function. However,
30 the inhibitory effects of aspirin and ticlopidine are dependent upon the agonist which activates the platelet. For example, aspirin is effective in blocking platelet aggregation induced by agonists such as collagen that are dependent upon the cyclooxygenase pathway. It is, however, less effective against concentrations of thrombin which can act by

- 2 -

cyclooxygenase independent pathways. Likewise, the inhibitory effects of ticlopidine, which inhibits ADP induced platelet aggregation, can be overcome by combinations of agonists. Thus, an efficacious platelet aggregation therapy that acts independently of the agonist and the pathway activating the platelet could be an important therapeutic advance giving greater efficacy than aspirin or ticlopidine alone in a broader spectrum of thrombotic events.

Integrin Superfamily

The firm attachment of endothelial cells to the subendothelial extracellular matrix is mediated via CAMs, which serve as receptors recognizing an array of adhesive proteins in the extracellular matrix. These proteins include von Willebrand factor (vWf), fibronectin, vitronectin, thrombospondin, laminins, collagen fibrils, elastin, microfibrils of elastin, and glycosaminoglycans. Most of the matrix adhesive molecules are the ligands for integrin receptors expressed in endothelial cells.

Integrins constitute an extended family ("superfamily") of membrane receptors interacting with adhesive proteins in plasma and extracellular matrix and with other membrane receptors (counter-receptors). The name "integrin" implies that they integrate the ligands on the outside of the cell with the cytoskeletal apparatus in the inside of the cell. Integrin receptors consist of a noncovalently linked Ca^{2+} -dependent, heterodimeric glycoprotein complex composed of α and β subunits. The eight known integrin β subunits give rise to eight families in which one "founder" β subunit forms heterodimers with different α subunits. There are at least 14 known α subunits. Receptors belonging to the β_1 and β_3 families are expressed in endothelial cells. The β_1 family, also named Very Late Antigens (VLA), is represented by the fibronectin receptor ($\alpha_5\beta_1$, or VLA-5), the collagen receptor ($\alpha_2\beta_1$, or VLA-2) and the laminin receptor ($\alpha_6\beta_1$). The β_3 family is represented by the vitronectin receptor ($\alpha_v\beta_3$), which is structurally similar (the same β_3 subunit) to the platelet integrin receptor for fibrinogen, glycoprotein (GP) IIb/IIIa complex (also referred to as $\alpha_{\text{IIb}}\beta_3$). The

- 3 -

functional difference between these two receptors is that the platelet receptor recognizes the γ chain domain (HHLGGAKQAGDV) of human fibrinogen and the endothelial vitronectin receptor does not. Both recognize the sequence R-G-D identified as the cell adhesion site of
5 fibrinectin, vitronectin, vWf, and the α chain of human fibrinogen. Therefore, synthetic peptides containing the R-G-D sequence cause detachment of endothelial cells from the extracellular matrix *in vitro*.

GP IIb/IIIa Antagonists

10 The final obligatory step in platelet aggregation is the binding of fibrinogen to an activated membrane-bound glycoprotein complex, GP IIb/IIIa. Platelet activators such as thrombin, collagen, epinephrine or ADP, are generated as an outgrowth of tissue damage. During activation, GP IIb/IIIa undergoes changes in conformation that
15 results in exposure of occult binding sites for fibrinogen. There are six putative recognition sites within fibrinogen for GP IIb/IIIa and thus fibrinogen can potentially act as a hexavalent ligand to crossing GP IIb/IIIa molecules on adjacent platelets. A deficiency in either fibrinogen or GP IIb/IIIa prevents normal platelet aggregation
20 regardless of the agonist used to activate the platelets. Since the binding of fibrinogen to its platelet receptor is an obligatory component of normal aggregation, GP IIb/IIIa is an attractive target for an antithrombotic agent.

Results from clinical trials of GP IIb/IIIa inhibitors support
25 this hypothesis. The monoclonal antibody 7E3, which blocks the GP IIb/IIIa receptor, has been shown to be an effective therapy for the high risk angioplasty population. It is used as an adjunct to percutaneous transluminal coronary angioplasty or atherectomy for the prevention of acute cardiac ischemic complications in patients at high risk for abrupt
30 closure of the treated coronary vessel. Although 7E3 blocks both the IIb/IIIa receptor and the $\alpha_v\beta_3$ receptor, its ability to inhibit platelet aggregation has been attributed to its function as a IIb/IIIa receptor binding inhibitor.

- 4 -

A study reported in The New England Journal of Medicine vol. 330, No. 14, pp. 956-961 (1994) showed a decrease from 12.8% to 8.3% in the combined endpoints of death, non-fatal myocardial infarction (MI) and need for urgent revascularization with fibrinogen receptor blockade. This benefit was at the expense of some additional risk of bleeding, with the need for transfusion increasing from 3% to 6%, and the incidence of patients with decreased hematocrit increasing from 7% to 15%. 7E3 was added to the standard regime of heparin and aspirin thus leaving few hemostatic control mechanisms intact. The clinical benefits of this drug could be seen at 6 months.

Many other studies have shown that blocking the GPIIb/IIIa receptor will stop platelet aggregation induced by all of the agonists and thus prevent thrombus formation but leave platelet adhesion relatively intact. The 7E3 monoclonal antibody is described in Coller *et al.*, *Ann. NY Acad. Sci.* 1991; 614:193-213; and Coller *et al.*, *J. Clin Invest.* 1985; 76:101-108. Others have used agents based on the RGD sequence, including snake venom proteins, small peptides, and peptidomimetics (Cook *et al.*, *Drugs of Future*, 1994; 19:135-159; and Cox *et al.*, *Medicinal Research Reviews*, 1994; 14:195-228).

The snake venom proteins, termed disintegrins, have provided important structural information, but their antigenicity has limited their development as therapeutic agents (Cook *et al.*, *ibid.*; and Cox *et al.*, *ibid.*). Integrilin (also known as INTEGRILIN™) is a cyclic peptide that is based on the KGD sequence in the snake venom protein barbourin (Cook *et al.*, *ibid.*; and Cox *et al.*, *ibid.*). It inhibits ligand binding to GP IIb/IIIa but has very little effect on ligand binding to $\alpha_v\beta_3$. Among the non-peptide compounds are Ro 44-9883 and MK-383, which are administered intravenously, and are also selective for GPIIb/IIIa (Cook *et al.*, *ibid.*; and Cox *et al.*, *ibid.*). Orally active agents include SC54684, which is a prodrug (i.e., it requires biotransformation *in vivo* to its active form) with high oral bioavailability and Ro 43-8857, GR144053, and DMP728, which are themselves the active inhibitors (Cook *et al.*, *ibid.*; and Cox *et al.*, *ibid.*). Literally thousands of other compounds have been synthesized in an

- 5 -

attempt to obtain optimal potency, metabolic stability, receptor specificity, and favorable intravascular survival. Despite variations in these compounds, virtually of all of them when they are in their active form retain the basic charge relations of the RGD sequence with a
5 positive charge separated from a negative charge by approximately 10-20 Å (Cook *et al.*, *ibid.*; and Cox *et al.*, *ibid.*).

Platelet aggregation is profoundly inhibited when increasing concentrations of murine 7E3 or c7E3 Fab are added to platelet-rich plasma *in vitro* or administered in incremental doses to
10 animals or humans *in vivo* (Coller *et al.*, *Ann. NY Acad.*, *ibid.*; Tchong *et al.*, *ibid.*; and Simoons *et al.*, *Circulation* 1994; 89:596-603). There is an excellent correlation between the percentage of receptors blocked and the inhibition of aggregation, with nearly complete inhibition of aggregation when 80% or more of the receptors are blocked (Coller *et al.*, *Ann. NY Acad.*, *ibid.*).
15

The results of the 7E3 study support the hypothesis that blockade of GPIIb/IIIa receptors is more effective than aspirin in preventing platelet thrombi, even in the presence of heparin. They also support the hypothesis that platelet-dependent thrombi frequently
20 contribute significantly to the development of ischemic complications after PTCA, even when minor mechanical dissections are present.

It has been clear for several decades that elevated blood cholesterol is a major risk factor for coronary heart disease, and many studies have shown that the risk of CHD events can be reduced by lipid-
25 lowering therapy. Prior to 1987, the lipid-lowering armamentarium was limited essentially to a low saturated fat and cholesterol diet, the bile acid sequestrants (cholestyramine and colestipol), nicotinic acid (niacin), the fibrates and probucol. Unfortunately, all of these treatments have limited efficacy or tolerability, or both. With the
30 introduction of lovastatin, the first inhibitor of HMG-CoA reductase to become available for prescription in 1987, for the first time physicians were able to obtain large reductions in plasma cholesterol with very few adverse effects.

- 6 -

Recent studies have unequivocally demonstrated that lovastatin, simvastatin and pravastatin, all members of the HMG-CoA reductase inhibitor class, slow the progression of atherosclerotic lesions in the coronary and carotid arteries. Simvastatin and pravastatin have
5 also been shown to reduce the risk of coronary heart disease events, and in the case of simvastatin a highly significant reduction in the risk of coronary death and total mortality has been shown by the Scandinavian Simvastatin Survival Study. This study also provided some evidence for a reduction in cerebrovascular events. Additional studies have shown
10 that HMG CoA RI's may have an effect on platelet aggregation.

Improved treatments for inhibiting platelet aggregation are currently being sought for the large number of individuals who are at risk for reocclusion following thrombolytic therapy and angioplasty, transient ischemic attacks and a variety of other vaso-occlusive
15 disorders. The instant invention addresses this problem by providing a combination therapy comprised of a platelet aggregation inhibitor with an HMG-CoA RI, and more particularly, a GP IIa/IIIb receptor antagonist with an HMG-CoA RI. When administered as part of a combination therapy, the platelet aggregation inhibitor together with the
20 HMG-CoA RI provide enhanced inhibition of platelet aggregation as compared to administration of either the HMG-CoA RI or the platelet aggregation inhibitor alone. Due to the greater benefit of the drug combination, lesser dosage amounts of the platelet aggregation inhibitor, and more particularly the GP IIa/IIIb receptor antagonist, may be
25 needed to achieve the desired clinical result, thereby resulting in improved safety.

SUMMARY OF THE INVENTION

One object of the instant invention is to provide a novel
30 combination therapy comprised of a therapeutically effective amount of a cholesterol reducing agent such as an HMG-CoA reductase inhibitor in combination with a platelet aggregation inhibitor which is useful for inhibiting platelet aggregation and for inhibiting the formation of thrombotic occlusions in mammals. The instant invention further

- 7 -

provides novel methods for treating, preventing and reducing the risk of occurrence of cardiovascular and cerebrovascular events and related vaso-occlusive disorders. Another object of this invention is to provide pharmaceutical compositions which can be used with the above-
5 described methods. A further object is to provide a kit comprised of an HMG-CoA reductase inhibitor composition and a platelet aggregation inhibitor composition. Additional objects will be evident from the following detailed description.

10 DETAILED DESCRIPTION OF THE INVENTION

The instant invention involves a novel combination therapy comprising the administration of a therapeutically effective amount of an HMG-CoA RI in combination with a therapeutically effective amount of a platelet aggregation inhibitor to a mammal, and more particularly,
15 to a human. The combination therapy is used to inhibit platelet aggregation in mammals who are in need of such inhibition, and to prevent or treat disorders related to platelet aggregation.

The instant invention also provides pharmaceutical compositions comprised of a therapeutically effective amount of an
20 HMG-CoA RI in combination with a therapeutically effective amount of a platelet aggregation inhibitor and a pharmaceutically acceptable carrier. One embodiment of the instant compositions is a single composition adapted for oral administration comprised of a therapeutically effective amount of an HMG-CoA RI in combination
25 with a therapeutically effective amount of a platelet aggregation inhibitor and a pharmaceutically acceptable carrier.

A compound which inhibits HMG-CoA reductase is used in combination with a platelet aggregation inhibitor to practice the instant invention. Examples of HMG-CoA reductase inhibitors that may be
30 used include but are not limited to lovastatin (MEVACOR®; see US Patent No. 4,231,938), simvastatin (ZOCOR®; see US Patent No. 4,444,784), pravastatin (PRAVACHOL®; see US Patent No. 4,346,227), fluvastatin (LESCOL®; see US Patent No. 5,354,772), atorvastatin (LIPITOR®; see US Patent No. 5,273,995) and cerivastatin

- 8 -

(also known as rivastatin; see US Patent No. 5,177,080). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996). Preferably, the HMG-CoA RI is selected from lovastatin and simvastatin.

The term HMG-CoA reductase inhibitor is intended to include all pharmaceutically acceptable salt, ester and lactone forms of compounds which have HMG-CoA reductase inhibitory activity, and therefore the use of such salts, esters and lactone forms is included within the scope of this invention.


Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the assays described or cited in U.S. Patent 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33.

A compound which inhibits platelet aggregation is used in combination with an HMG-CoA reductase inhibitor to practice the instant invention. In one embodiment of the instant invention, the compound which inhibits platelet aggregation is an antagonist for the glycoprotein IIb/IIIa fibrinogen receptor. Examples of glycoprotein IIb/IIIa receptor antagonists are described in United States Patent No.'s 5,470,849, 5,463,011, 5,455,243, 5,451,578, 5,446,056, 5,441,952, 5,422,249, 5,416,099, 5,405,854, 5,397,791, 5,393,760, 5,389,631, 5,380,713, 5,374,622, 5,358,956, 5,344,783, 5,340,798, 5,338,723, 5,334,596, 5,321,034, 5,318,899 (e.g. cyclic heptapeptides such as Mpr-(Acetimidyl-Lys)-Gly-Asp-Trp-Phe-Cys-NH₂, Mpr-(Acetimidyl-Lys)-Gly-Asp-Trp-Phe-Pen-NH₂, Mpr-(Phenylimidyl-Lys)-Gly-Asp-Trp-Phe-Pen-NH₂, and Mpr-(Phenylimidyl-Lys)-Gly-Asp-Trp-Phe-Cys-NH₂, wherein Mpr is mercapto propionyl), 5,312,923, 5,294,616, 5,292,756, 5,281,585, 5,272,158, 5,264,420, 5,260,307, 5,239,113 (e.g. ethyl 3-[[4-[[4-(aminoiminomethyl)phenyl]amino]-1,4-dioxobutyl]amino]-4-pentynoate), 5,227,490, 5,206,373, 4,703,036 (e.g. N-Methyl-D-phenylalanyl-N-[(1S)-1-formyl-4-guanidinobutyl]-L-prolinamide); European Patent publication No.'s EP 505 868 (e.g., ((1-

- 9 -

(2-((4-(aminoiminomethyl)benzoyl)amino)-3-(4-hydroxyphenyl)-1-oxopropyl)-4-piperidinyloxy)-(S)-acetic acid), EP 333 356, and EP 656 348; and International Publication No.'s WO 93/11152 (e.g., N-(2-(2-
 5 (((3-((aminoiminomethyl)amino)propyl)amino)-carbonyl)-1-piperidnyl)-1-(cyclohexylmethyl)-2-oxoethyl)-(R,S)-glycine), WO 94/18981, WO 94/22820, WO 95/14683, and WO 97/15568, all of which are herein incorporated by reference, and wherein the scope of this invention includes, but is not limited to, the use of each of the specifically disclosed compounds therein. They are described as
 10 useful for inhibiting fibrinogen binding and inhibiting clot formation.

In particular, the GP IIb/IIIa receptor antagonist is selected from the following compounds and the pharmaceutically acceptable salts, esters, and solvates (including hydrates) thereof: [3(R)-[2-piperidin-4-yl)ethyl]-2-piperidone-1]acetyl-3(R)-methyl-β-alanine
 15 described in United States Patent No. 5,281,585, (see compound 57 in column 67) and referred to herein as Compound A:




Compound A;


20

5-[(4-Piperidinyloxy)methoxy]-2-indolecarbonyl-2(S)-phenylsulfonyl-amino-β-alanine described in WO 97/15568 at page 20 as compound 2-6, and referred to herein as Compound B:

- 10 -


Compound B;

2(S)-[(p-Toluenesulfonyl)amino]-3-[[[5,6,7,8-tetrahydro-4-oxo-5-[2-(piperidin-4-yl)ethyl]-4H-pyrazolo-[1,5-a][1,4]diazepin-2-yl]carbonyl]-amino]propionic acid described in WO 94/18981, and referred to herein as Compound C:

Compound C;


MK-383 (2-S-(n-Butylsulfonylamino)-3[4-piperidin-4-yl]butyloxyphenyl]propionic acid hydrochloride, and also known as tirofiban) described in United States Patent 5,292,756; DMP 728; DMP 754 ((R)-methyl-3-[[[3-[4-(aminoiminomethyl)phenyl]-4,5-dihydro-5-isoxazolyl]acetyl]amino]-N-(butoxycarbonyl)-L-alanine monoacetate) from DuPont-Merck, described in WO 95/14683 and in *Tetrahedron Letters*, 1996, 37 :4455-4458:

- 11 -




- 5 Ro44-9883, Ro43-8857 and Ro48-3657 (acetic acid, [[1-[2-[[4-[amino(hydroxyimino)methyl]benzoyl]amino]-1-oxopropyl]-4-piperidinyl]oxy]-, ethyl ester, and also known as sibrafiban) from Hoffman-LaRoche; sibrafiban and related compounds are described in EP 656 348:

10



- 15 xemlofiban (ethyl 3-[[4-[[4-(aminoiminomethyl)phenyl]amino]-1,4-dioxobutyl]amino]-4-pentynoate, also known as xemilofiban and SC-54684), particularly the HCl salt thereof, described in U.S. Patent No.'s 5,344,957 and 5,239,113:




20

- 12 -


fradafiban ((3S,5S)-5-[[[4'-Amidino-4-biphenyl)oxy]methyl]-2-oxo-3-pyrrolidineacetic acid, also known as BIBU-104) as described in U.S. Patent No. 5,541,343 assigned to Thomae:

5



Fradafiban ;

orbofiban (N-[[[(3S)-1-(p-Amidinophenyl)-2-oxo-3-pyrrolidinyl-carbamoyl]-β-alanine, ethyl ester), particularly the monoacetate and
10 monoacetate hydrate forms thereof, as described in U.S. Patent No. 5,484,946 assigned to Searle:



Orbofiban ;

15

SB 214857 ((-)-(S)-2-[7-(4,4'-Bipiperidin-1-ylcarbonyl)-4-methyl-3-oxo-2,3,4,5-tetrahydro-1H-1,4-benzodiazepin-2-yl]acetic acid) from SmithKline Beecham, as described in WO 95/18619; ZD-2486 ((R)-3-Methyl-4-[4-[4-(4-pyridyl)piperazin-1-yl]phenoxy]butyric acid) from
20 Zeneca, as described in U.S. Patent No.'s 5,556,977 and 5,563,141; TAK-029 from Takeda; RPR 109891 from Rhone Polenc Rorer; GR144053 from Glaxo; GR233548 from Glaxo; and SDZ 562 from Sandoz.

The compounds MK-383, DMP 728, Ro44-9883, Ro43-
25 8857, SC-54684 and GR144053 are described in Cook *et al.*, *Drugs of*

- 13 -

the Future, 1994, 19(2) :135-159, and Cox *et al.*, *Medicinal Research Reviews*, 1994, 14 :195-228. DMP 728 is also described in *Circulation*, 1996, 93 :537-543; and GR144053 is also described in *Thrombosis and Hematosis*, 1993, 69 :1071. TAK 029 is described in *J. Pharmacology and Experimental Therapeutics*, 1996, 277 :502-510. Xemlofiban is
5 described in *Circulation*, 1995, 92 :2331.

More particularly, the GP IIb/IIIa receptor antagonist is selected from Compound A, Compound B, and DMP 754, which are all orally available compounds. Most particularly, the GP IIb/IIIa receptor
10 antagonist is DMP 754.

One test which is used to evaluate fibrinogen IIb/IIIa receptor antagonist activity is based on evaluation of inhibition of ADP-stimulated platelets. Aggregation requires that fibrinogen bind to and occupy the platelet fibrinogen receptor site. Inhibitors of fibrinogen
15 binding inhibit aggregation. In the ADP-stimulated platelet aggregation assay, human platelets are isolated from fresh blood, collected into acid citrate/dextrose by differential centrifugation followed by gel filtration on Sepharose 2B in divalent ion-free Tyrode's buffer (pH 7.4) containing 2% bovine serum albumin.

20 Platelet aggregation is measured at 37°C in a Chronolog aggregometer. The reaction mixture contains gel-filtered human platelets (2×10^8 per ml), fibrinogen (100 micrograms per ml (ug/ml)), Ca^{2+} (1 mM), and the compound to be tested. The aggregation is initiated by adding 10 mM ADP 1 minute after the other components
25 are added. The reaction is then allowed to proceed for at least 2 minutes. The extent of inhibition of aggregation is expressed as the percentage of the rate of aggregation observed in the absence of inhibitor. The IC_{50} is the dose of a particular compound inhibiting aggregation by 50% relative to a control lacking the compound.

30 Compounds which are selective GP IIb/IIIa receptor antagonists may be employed in the instant invention, as well as compounds which block both the GP IIb/IIIa receptor and the $\alpha_v\beta_3$ vitronectin receptor, such as the monoclonal antibody 7E3. Compounds which are selective for the glycoprotein IIb/IIIa receptor are those

- 14 -

having a preference (e.g. 10-fold) for binding to IIb/IIIa as compared to other receptors of the integrin family (e.g. $\alpha_v\beta_3$, $\alpha_5\beta_1$, $\alpha_v\beta_5$).

Selectivity of these compounds can be readily determined by persons skilled in the art.

5 In another embodiment of this invention, the compound which inhibits platelet aggregation is one that blocks ADP induced platelet aggregation. Examples of such compounds include ticlopidine (TICLID®), and clopidogrel (PLAVIX®).

10 In a further embodiment of this invention, the compound which inhibits platelet aggregation is selected from aspirin and dipyridamole.

 Herein, the term platelet aggregation inhibitor (or inhibitor of platelet aggregation) is intended to include all pharmaceutically acceptable salt, ester and solvate forms, including hydrates, of
15 compounds which have platelet aggregation inhibitory activity as well as pro-drug forms. Such pro-drugs are compounds which do not have platelet aggregation inhibitory activity outside the body but become active as inhibitors after they are administered to the patient. Therefore the use of such salts, esters solvate forms and pro-drugs of platelet
20 aggregation inhibitors is included within the scope of this invention.

 Likewise, the term GP IIb/IIIa receptor antagonist is intended to include all pharmaceutically acceptable salt, ester and solvate forms, including hydrates, of compounds which have GP IIb/IIIa receptor antagonist activity as well as pro-drug forms. Such pro-drugs
25 are compounds which do not have GP IIb/IIIa receptor antagonist activity outside the body but become active as antagonists after they are administered to the patient. Therefore the use of such salts, esters, solvate forms and pro-drugs of GP IIb/IIIa receptor antagonists is also included within the scope of this invention. Pro-drug forms of IIb/IIIa
30 receptor antagonists generally are not active antagonists until after they are metabolised in the body to the active drug form; such prodrugs may be, but are not limited to, ester derivatives. Ester derivatives of the described compounds may act as prodrugs which, when absorbed into the bloodstream of a warm-blooded animal, may cleave in such a

- 15 -

manner as to release the drug form and permit the drug to afford improved therapeutic efficacy. An example of such a pro-drug is Ro 48-3657.

Herein, the term "pharmaceutically acceptable salts" shall
5 mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base. Examples of such salts include, but are not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride,
10 clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycollylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynapthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate,
15 methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamaote, palmitate, panthothenate, phosphate/diphosphate, polygalacturonate, salicylate, sodium, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide, valerate.

The terms "active agent(s)" and "active drug(s)" are used
20 herein as a way to refer to both the HMG-CoA reductase inhibitors and the platelet aggregation inhibitors which are employed in the instant methods and compositions. Both terms are intended to encompass all salt, ester and pro-drug forms of HMG-CoA reductase inhibitors and platelet aggregation inhibitors, even where the pro-drug form is not
25 active itself but is converted to the active drug form after administration.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients, in specified amounts where amounts are specified, as well as any product which
30 results directly or indirectly from combination of the specified ingredients, in the specified amounts where amounts are specified.

As used herein, the term "myocardial infarction" is intended to include both Q-wave and non-Q-wave myocardial infarction, unless otherwise noted.

- 16 -

The instant method involves the administration of an HMG-CoA reductase inhibitor in combination with a platelet aggregation inhibitor. This combination therapy includes administration of a single pharmaceutical dosage formulation which contains both the HMG-CoA reductase inhibitor and the platelet aggregation inhibitor, as well as administration of each active agent in its own separate pharmaceutical dosage formulation. While the HMG-CoA reductase inhibitor can be administered orally or parenterally, oral dosing is preferred. The GP IIb/IIIa receptor antagonist can be administered orally, intravenously, transdermally (for example using an iontophoretic patch), intraocularly, intranasally or by other routes known to those skilled in the medical arts, taking into account that certain GP IIb/IIIa receptor antagonists are developed for oral administration while others may be developed for non-oral routes such as intravenous administration. Ticlopidine, clopidogrel, aspirin and dipyridamole are administered orally. Preferably, both active agents of the instant combination therapy are administered orally, and most preferably the active agents are combined in a single oral dosage formulation.

For example, a GP IIb/IIIa receptor antagonist and an HMG-CoA reductase inhibitor can be administered to the patient together in one oral composition such as a tablet or capsule. Another example would be a single oral composition comprised of aspirin and an HMG-CoA reductase inhibitor. Alternatively, the combination therapy may comprise administration of an oral HMG-CoA reductase inhibitor composition with a separate oral aspirin composition, or with a separate GP IIb/IIIa receptor antagonist composition formulated for oral, intravenous, transdermal, intraocular, or intranasal administration.

Where separate dosage formulations are used, the HMG-CoA reductase inhibitor and the platelet aggregation inhibitor can be administered at essentially the same time, i.e., concurrently, or at separately staggered times, i.e., sequentially. Similarly, a GP IIb/IIIa receptor antagonist may be administered on an intravenous regimen, while the patient is orally dosed once a day with a conventional or controlled-release formulation of an HMG-CoA reductase inhibitor.

- 17 -

Combination therapy is understood to include all these regimens. Administration in these various ways are suitable for the present invention as long as the beneficial pharmaceutical effect of the platelet aggregation inhibitor and HMG-CoA reductase inhibitor are realized by the patient at substantially the same time. Such beneficial effect is
5 achieved when the target blood level concentrations of each active drug are maintained at substantially the same time. Concurrent oral administration is preferred.

It is expected that a combination therapy of intravenously administered GP IIb/IIIa receptor antagonist with orally administered
10 HMG-CoA reductase inhibitor could be used in response to an acute medical event where inhibition of platelet aggregation is needed, and may generally be administered for a period of time of one or two weeks or up to a month or longer if deemed necessary. Where the
15 combination therapy involves for example oral administration of both the GP IIb/IIIa receptor antagonist and the HMG-CoA reductase inhibitor, the therapy may be administered on a longer-term chronic basis, such as a period of several months or years, for as long as deemed medically appropriate for the patient.

20 Therapeutically effective amounts of the platelet aggregation inhibitors and the HMG-CoA reductase inhibitors are suitable for use in the compositions and methods of the present invention. The term "therapeutically effective amount" is intended to mean that amount of a drug or pharmaceutical agent that will elicit the
25 biological or medical response of a tissue, a system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician. The dosage regimen utilizing an HMG-CoA RI in combination with a platelet aggregation inhibitor is selected in accordance with a variety of factors including type, species, age, weight,
30 sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt or ester thereof employed. Since two different active agents are being used together in a combination therapy, the potency of each of the agents and the enhanced

- 18 -

effects achieved by combining them together must also be taken into account. A consideration of these factors is well within the purview of the ordinarily skilled clinician for the purpose of determining the therapeutically effective amounts of the drug combination needed to prevent, counter, or arrest the progress of the condition.

Dosage information for HMG-CoA RI's is well known in the art, since several HMG-CoA RI's are marketed in the U.S. In particular, the daily dosage amounts of the HMG-CoA reductase inhibitor may be the same or similar to those amounts which are employed for anti-hypercholesterolemic treatment and which are described in the *Physicians' Desk Reference* (PDR). For example, see the 50th Ed. of the PDR, 1996 (Medical Economics Co); in particular, see at page 216 the heading "Hypolipidemics," sub-heading "HMG-CoA Reductase Inhibitors," and the reference pages cited therein.

Preferably, the oral dosage amount of HMG-CoA RI is from about 1 to 200 mg/day, and more preferably from about 5 to 160 mg/day. However, dosage amounts will vary depending on the potency of the specific HMG-CoA reductase inhibitor used as well as other factors as noted above. Also, the dosage amount of HMG-CoA reductase inhibitor needed to achieve the desired effect will be affected by the dosage amount and potency of the IIb/IIIa receptor antagonist or other platelet aggregation inhibitor which is employed in the combined therapy. An HMG-CoA RI which has sufficiently greater potency may be given in sub-milligram daily dosages.

As examples, the daily dosage amount for simvastatin may be selected from 5 mg, 10 mg, 20 mg, 40 mg, 80 mg and 160 mg; for lovastatin, 10 mg, 20 mg, 40 mg and 80 mg; for fluvastatin sodium, 20 mg, 40 mg and 80 mg; and for pravastatin sodium, 10 mg, 20 mg, and 40 mg. The daily dosage amount for atorvastatin calcium may be in the range of from 1 mg to 160 mg, and more particularly from 5 mg to 80 mg, including dosage amounts of 10 mg, 20 mg and 40 mg. Oral administration may be in single or divided doses of two, three, or four times daily, although a single daily dose of the HMG-CoA RI is preferred.

- 19 -

Oral dosages of GP IIb/IIIa receptor antagonists when used for the indicated effects, will range between about 0.001 mg per kg of body weight per day (mg/kg/day) to about 50 mg/kg/day and preferably 0.005-20 mg/kg/day and most preferably 0.005-10 mg/kg/day. Suitable oral tablets and capsules contain between 0.1 mg and 5 g, preferably between 0.5 mg and 2 g, most preferably between 0.5 mg and 1 g, for example, 0.5 mg, 1 mg, 5 mg, 10 mg, 50 mg, 150 mg, 250 mg, or 500 mg of GP IIb/IIIa receptor antagonist. Oral administration may be in one or divided doses of two, three, or four times daily. A single daily dose is preferred.

Intravenously, the most preferred doses for GP IIb/IIIa receptor antagonists will range from about 0.5 μ g to about 5 mg/kg/minute during a constant rate infusion, to achieve a plasma level concentration during the period of time of administration of between 0.1 ng/ml and 1 μ g/ml.

Dosage amounts for ticlopidine are described in the *Physicians' Desk Reference*. Dosage amounts of aspirin for the indicated effects are known to those skilled in medical arts, and generally range from about 75 mg to about 325 mg per day. For example, a formulation may contain 75 mg, 80 mg, 160 mg, 250 mg, or 325 mg of aspirin.

Suitable oral formulations of clopidogrel may contain from 25 mg to 500 mg, preferably from 75 mg to 375 mg, and most preferably from 75 mg to 150 mg of clopidogrel. For example, the formulation may contain 25 mg, 50 mg, 75 mg, 150 mg, 250 mg, or 500 mg of clopidogrel. Oral administration may be in one or divided doses of two, three, or four times daily. A single daily dose is preferred.

The active agents employed in the instant combination therapy can be administered in such oral forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. The instant invention includes the use of both oral rapid-release and time-controlled release pharmaceutical formulations (see, e.g., U.S.

- 20 -

Patent No. 5,366,738 which describes controlled release formulations). Suitable intravenous compositions for the GP IIb/IIIa receptor antagonists include bolus or extended infusion. Such oral and intravenous compositions are known to those of ordinary skill in the pharmaceutical arts. For example, see *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA.

The active drugs can be administered in admixture with pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with a non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, modified sugars, modified starches, methyl cellulose and its derivatives, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and other reducing and non-reducing sugars, magnesium stearate, steric acid, sodium stearyl fumarate, glyceryl behenate, calcium stearate and the like. For oral administration in liquid form, the drug components can be combined with non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring and flavoring agents can also be incorporated into the mixture. Stabilizing agents such as antioxidants (BHA, BHT, propyl gallate, sodium ascorbate, citric acid) can also be added to stabilize the dosage forms. Other suitable components include gelatin, sweeteners, natural and synthetic gums such as acacia, tragacanth or alginates, carboxymethylcellulose, polyethylene glycol, waxes and the like.

The active drugs can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be

- 21 -

formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Active drug may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. Active drug may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxy-ethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, active drug may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels.

The compositions and methods of the present invention may be used as treatment for acute cardiovascular and cerebrovascular events, as well as for chronic therapy for prevention or reduction of risk of occurrence of cardiovascular and cerebrovascular events. For example, the compositions of this invention, and methods for administering the combination therapy of a platelet aggregation inhibitor with an HMG-CoA reductase inhibitor, are useful for treating, preventing or reducing the risk of occurrence of acute coronary ischemic syndrome in mammals, and more particularly in humans, who are at risk of developing acute coronary ischemic syndrome. Acute coronary ischemic syndrome includes the conditions of unstable angina and non-Q-wave myocardial infarction.

Compositions and methods of the invention may be used to prevent or reduce the risk of formation of thrombi and thromboemboli and therefore to prevent or reduce the risk of thrombotic occlusions and reocclusions. They are useful in surgery on peripheral arteries (arterial grafts, carotid endarterectomy) and in cardiovascular surgery where manipulation of arteries and organs, and/or the interaction of platelets with artificial surfaces, leads to platelet aggregation and potential

- 22 -

formation of thrombi and thromboemboli. For example, the combination therapy can be used for preventing or reducing the risk of occurrence of platelet thrombosis, thromboembolism and reocclusion after acute intervention such as atherectomy, angioplasty (PTCA), coronary artery bypass procedures or cardiac valve replacement. The combination therapy can also be used for preventing or reducing the risk of occurrence of platelet thrombosis, thromboembolism and reocclusion during and after thrombolytic therapy. Since blood vessels can also sustain chronic damage by the pathophysiological processes of atherosclerosis, patients with atherosclerosis can also be treated with the instant combination therapy to prevent or reduce the risk of occlusive thrombus formation. The instant combination therapy can be used to treat, prevent or reduce the risk of intermittent claudication, which is a clinical manifestation of peripheral vessel disease. Combination therapy of an HMG-CoA reductase inhibitor with a GP IIb/IIIa receptor antagonist may reduce the risk of thrombocytopenia.

The instant combination therapy can also be used to treat, prevent or reduce the risk of a first or subsequent Q-wave myocardial infarction in persons at risk for such events as well as to prevent or reduce the risk of restenosis in persons at risk for restenosis. Additionally, the instant combination therapy can be used for treating, preventing or reducing the risk of occurrence of acute cerebrovascular ischemic events such as a first or subsequent thrombotic stroke or transient ischemic attack. In general, the instant combination therapy can be used whenever antiplatelet therapy, or inhibition of platelet aggregation, is needed.

The compositions and methods of the present invention are also useful in combination with procedures for treating patients with other anticoagulants (e.g. thrombin inhibitors such as heparin and Factor Xa inhibitors such as warfarin), and thrombolytic agents (e.g. streptokinase and tissue plasminogen activator). The instant combination therapy can also be co-administered with a β -adrenergic receptor blocker. In particular, a combination of an HMG-CoA RI with aspirin can be co-administered with a β -adrenergic receptor

- 23 -

blocker to reduce the risk of coronary heart disease and cerebrovascular clinical events such as myocardial infarction, stroke and cardiovascular death, particularly in post-MI patients. Examples of β -adrenergic receptor blockers include but are not limited to
5 acebutolol, atenolol, betaxolol, bioprolol, carteolol, labetalol, metoprolol, nadolol, penbutolol, pindolol, propanolol, and timolol. Dosage amounts of β -adrenergic receptor blockers are described in the PDR.

 In accordance with this invention, a therapeutically
10 effective amount of an HMG-CoA RI and a therapeutically effective amount of a platelet aggregation inhibitor can be used for the preparation of a medicament useful for inhibiting platelet aggregation, and for treating, preventing or reducing the risk of developing acute coronary ischemic syndrome in mammals, particularly in humans.
15 Additionally, a therapeutically effective amount of an HMG-CoA RI and a therapeutically effective amount of a platelet aggregation inhibitor can be used for the preparation of a medicament useful for preventing or reducing the risk of formation of thrombi and thromboemboli, for preventing or reducing the risk of thrombotic occlusions and
20 reocclusions, for treating, preventing or reducing the risk of a first or subsequent myocardial infarction, for preventing or reducing the risk of restenosis, for treating, preventing or reducing the risk of acute cerebrovascular ischemic events such as a first or subsequent thrombotic stroke or transient ischemic attack, and for halting or slowing the
25 progression of atherosclerotic disease. More particularly, a therapeutically effective amount of an HMG-CoA RI and a therapeutically effective amount of a platelet aggregation inhibitor can be used together for the preparation of a medicament suitable for oral administration which is useful for the above-described treatments.
30 Similarly, a therapeutically effective amount of an HMG-CoA RI can be used for the preparation of a medicament for use in combination with a therapeutically effective amount of a platelet aggregation inhibitor, which is useful for the above-described treatments. Also, a therapeutically effective amount of a platelet aggregation inhibitor can

- 24 -

be used for the preparation of a medicament for use in combination with a therapeutically effective amount of an HMG-CoA RI, which is useful for the above-described treatments.

An additional embodiment of the instant invention
5 involves a kit comprised of an HMG-CoA RI in an oral dosage formulation and a platelet aggregation inhibitor in a separate oral dosage formulation. More particularly, the kit is comprised of an HMG-CoA RI selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin; and
10 the platelet aggregation inhibitor is selected from the group consisting of a GP IIb/IIIa receptor antagonist, ticlopidine, clopidogrel, aspirin and dipyridamole. In one class of this embodiment the HMG-CoA RI is selected from lovastatin and simvastatin, and more particularly the HMG-CoA RI is simvastatin.
15 In a second class of this embodiment, the platelet aggregation inhibitor is a GP IIb/IIIa receptor antagonist selected from the group consisting of Compound A, Compound B, and DMP 754. In a third class of this embodiment, the platelet aggregation inhibitor is aspirin.

One example of this embodiment is a kit comprised of
20 an oral dosage formulation of simvastatin and an oral dosage formulation of aspirin. The packaging for the kit could be designed and manufactured in a variety of ways. A preferred example is a blister package containing rows of a simvastatin tablet and an aspirin tablet placed side by side on the same blister card, each of the two
25 tablets in its own blister bubble, with calendar or similar type markings on the card that convey to the user that one "pair" of tablets (i.e., one simvastatin tablet and one aspirin tablet) is to be ingested per day. A kit containing simvastatin and a GP IIb/IIIa receptor antagonist such as Compound A, Compound B, or DMP 754
30 could be designed in a similar fashion.

A further class of this embodiment involves the kit described above further comprising an oral dosage formulation of a pharmaceutically active agent selected from a β -adrenergic receptor blocker and an angiotensin II receptor antagonist, in addition to the

- 25 -

HMG-CoA RI and the platelet aggregation inhibitor. More particularly, β -adrenergic receptor blockers without intrinsic sympathomimetic activity (ISA) and without alpha blocking properties have a cardioprotective effect for patients who have had a myocardial infarction and can be employed in the kit. Atenolol, metoprolol, betaxolol and acebutolol are beta blockers without ISA. Losartan potassium is currently marketed in the U.S. under the trademark COZAAR®, and is one example of an angiotensin II receptor antagonist that could be employed in the kit. Dosages for beta blockers and for losartan potassium are commonly known to those skilled in the pharmaceutical arts and can be found, for example in the *Physician's Desk Reference*.

Examples of dosage formulations are as follows.

- 26 -

EXAMPLE 1
Tablet Preparation

Tablets containing simvastatin and a glycoprotein
5 IIb/IIIa receptor antagonist can be prepared as illustrated below:

<u>Ingredient</u>	<u>Amount</u>
simvastatin	1 mg - 200 mg.
glycoprotein IIb/IIIa receptor antagonist	0.1 mg - 5 g
10	
diluent	} excipients qs. 200 - 400 mg.
binder	
disintegrant	
lubricant	

EXAMPLE 2
Tablet Preparation

15

Tablets containing 25.0, 50.0, and 100.0 mg, respectively,
of a GP IIb/IIIa receptor antagonist are prepared as illustrated below:

20 TABLE FOR DOSES CONTAINING
FROM 25-100MG OF GP IIb/IIIa RECEPTOR ANTAGONIST

	<u>Amount-mg</u>		
GP IIb/IIIa receptor antagonist	25.0	50.0	100.0
Microcrystalline cellulose	37.25	100.0	200.0
Modified food corn starch	37.25	4.25	8.5
Magnesium stearate	0.50	0.75	1.5

- 27 -

All of the active compound, cellulose, and a portion of the corn starch are mixed and granulated to 10% corn starch paste. The resulting granulation is sieved, dried and blended with the remainder of the corn starch and the magnesium stearate. The resulting granulation is then compressed into tablets containing 25.0, 50.0, and 100.0 mg, respectively, of active ingredient per tablet.

EXAMPLE 3

Intravenous formulations

10

An intravenous dosage form of the GP IIb/IIIa receptor antagonist is prepared as follows:

	<u>Amount</u>
GP IIb/IIIa receptor antagonist	0.5-10.0mg
Sodium Citrate	5-50mg
Citric Acid	1-15mg
Sodium Chloride	1-8mg
Water for Injection (USP)	q.s. to 1 L

15

Utilizing the above quantities, the active compound is dissolved at room temperature in a previously prepared solution of sodium chloride, citric acid, and sodium citrate in Water for Injection (USP, see page 1636 of United States Pharmacopeia/National Formulary for 1995, published by United States Pharmacopeial Convention, Inc., Rockville, Maryland, copyright 1994.

20

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the particular dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications for the active agents

25

- 28 -

used in the instant invention as indicated above. Likewise, the specific pharmacological responses observed may vary according to and depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type
5 of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as
10 broadly as is reasonable.

- 29 -

WHAT IS CLAIMED IS:

1. A method for reducing the risk of developing a condition selected from the group consisting of:
 - 5 acute coronary ischemic syndrome, thrombosis, thromboembolism, thrombotic occlusion and reocclusion, restenosis, first or subsequent Q-wave myocardial infarction, transient ischemic attack, and first or subsequent thrombotic stroke comprising administration of a therapeutically effective amount of an HMG-CoA reductase inhibitor in
 - 10 combination with a therapeutically effective amount of a platelet aggregation inhibitor to a mammal at risk of developing one or more of said conditions.
2. The method of Claim 1 wherein the platelet aggregation inhibitor is a glycoprotein IIb/IIIa receptor antagonist.
 - 15
3. The method of Claim 1 wherein the platelet aggregation inhibitor is selected from the group consisting of ticlopidine and clopidogrel.
 - 20
4. The method of Claim 1 wherein the platelet aggregation inhibitor is selected from the group consisting of aspirin and dipyridamole.
 - 25
5. The method of Claim 1 wherein the condition is acute coronary ischemic syndrome.
 - 30
6. The method of Claim 1 wherein the condition is selected from thrombosis, thromboembolism, thrombotic occlusion and reocclusion.
 - 35
7. The method of Claim 1 wherein the condition is restenosis.

- 30 -

8. The method of Claim 1 wherein the condition is Q-wave myocardial infarction.
9. The method of Claim 1 wherein the condition is
5 transient ischemic attack.
10. The method of Claim 1 wherein the condition is thrombotic stroke .
- 10 11. A method for treating a cardiovascular event selected from acute coronary ischemic syndrome and Q-wave myocardial infarction comprising administration of a therapeutically effective amount of an HMG-CoA reductase inhibitor in combination with a therapeutically effective amount of a platelet aggregation inhibitor to a
15 mammal in need of such treatment.
12. A method for treating a cerebrovascular event selected from thrombotic stroke and transient ischemic attack comprising administration of a therapeutically effective amount of an
20 HMG-CoA reductase inhibitor in combination with a therapeutically effective amount of a platelet aggregation inhibitor to a mammal in need of such treatment.
13. The method of Claim 11 wherein the platelet
25 aggregation inhibitor is selected from the group consisting of a glycoprotein IIb/IIIa receptor antagonist, ticlopidine, clopidogrel, aspirin and dipyridamole.
14. The method of Claim 12 wherein the platelet
30 aggregation inhibitor is selected from the group consisting of a glycoprotein IIb/IIIa receptor antagonist, ticlopidine, clopidogrel, aspirin and dipyridamole.

- 31 -

15 15. A method for inhibiting platelet aggregation
 comprising administration of a therapeutically effective amount of a
 platelet aggregation inhibitor in combination with a therapeutically
 effective amount of an HMG-CoA reductase inhibitor to a mammal in
 need of such treatment.

10 16. The method of Claim 15 wherein the platelet
 aggregation inhibitor is selected from the group consisting of a
 glycoprotein IIb/IIIa receptor antagonist, ticlopidine, clopidogrel,
 aspirin and dipyridamole.

15 17. The method of Claim 2 wherein the glycoprotein
 IIb/IIIa receptor antagonist and the HMG-CoA reductase inhibitor are
 both administered orally.

18. The method of Claim 2 wherein the glycoprotein
 IIb/IIIa receptor antagonist and the HMG-CoA reductase inhibitor are
 administered in a single oral dosage composition.

20 19. The method of Claim 2 wherein the glycoprotein
 IIb/IIIa receptor antagonist is administered intravenously and the HMG-
 CoA reductase inhibitor is administered orally.

25 20. The method of Claim 2 wherein the glycoprotein
 IIb/IIIa receptor antagonist is selected from the group consisting of:

 Mpr-(Acetimidyl-Lys)-Gly-Asp-Trp-Phe-Cys-NH₂,
 Mpr-(Acetimidyl-Lys)-Gly-Asp-Trp-Phe-Pen-NH₂,
 Mpr-(Phenylimidyl-Lys)-Gly-Asp-Trp-Phe-Pen-NH₂,
30 Mpr-(Phenylimidyl-Lys)-Gly-Asp-Trp-Phe-Cys-NH₂,
 N-Methyl-D-phenylalanyl-N-[(1S)-1-formyl-4-guanidinobutyl]-L-
 prolinamide,
 ((1-(2-((4-(aminoiminomethyl)benzoyl)amino)-3-(4-hydroxyphenyl)-1-
 oxopropyl)-4-piperidinyl)oxy)-(S)-acetic acid,

- 32 -

- N-(2-(2-(((3-((aminoiminomethyl)amino)propyl)amino)carbonyl)-1-piperidnyl)-1-(cyclohexylmethyl)-2-oxoethyl)-(R,S)-glycine, Ethyl 3-[[4-[[4-(aminoiminomethyl)phenyl]amino]-1,4-dioxobutyl]amino]-4-pentynoate,
- 5 (2-S-(n-Butylsulfonylamino)-3[4-(piperidin-4-yl)butyloxyphenyl]propionic acid hydrochloride, and 2(S)-[(p-Toluenesulfonyl)amino]-3-[[[5,6,7,8-tetrahydro-4-oxo-5-[2-(piperidin-4-yl)ethyl]-4H-pyrazolo-[1,5-a][1,4]diazepin-2-yl]carbonyl]-amino]propionic acid;
- 10 and the pharmaceutically acceptable salts, esters and pro-drugs thereof.

21. The method of Claim 2 wherein the glycoprotein IIb/IIIa receptor antagonist is selected from the group consisting of: DMP 754, [3(R)-[2-piperidin-4-yl)ethyl]-2-piperidone-1]acetyl-3(R)-methyl- β -alanine, and 5-[(4-Piperidinyl)methoxy]-2-indolecarbonyl-2(S)-phenylsulfonyl-amino- β -alanine and the pharmaceutically acceptable salts, esters and pro-drugs thereof.

15

22. The method of Claim 21 wherein the glycoprotein IIb/IIIa receptor antagonist is DMP754 and the pharmaceutically acceptable salts, esters and pro-drugs thereof.

20

23. The method of Claim 1 wherein the HMG-CoA reductase inhibitor is selected from the group consisting of lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin and cerivastatin and the pharmaceutically acceptable lactone, salt and ester forms thereof.

25

24. The method of Claim 23 wherein the HMG-CoA reductase inhibitor is selected from the group consisting of lovastatin and simvastatin and the pharmaceutically acceptable salts and esters thereof.

30

25. A method for preventing or reducing the risk of occurrence of acute coronary ischemic syndrome and acute

- 33 -

cerebrovascular ischemic events in patients at risk for said occurrences comprising inhibiting the binding of fibrinogen to the glycoprotein IIb/IIIa receptor and inhibiting the activity of HMG-CoA reductase.

5 26. A pharmaceutical composition comprising a platelet aggregation inhibitor, an HMG-CoA reductase inhibitor and a pharmaceutically acceptable carrier.

10 27. The composition of Claim 26 wherein the HMG-CoA reductase inhibitor is selected from lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, cerivastatin and the pharmaceutically acceptable lactone, salt and ester forms thereof.

15 28. The composition of Claim 27 wherein the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin and the pharmaceutically acceptable salts and esters thereof.

20 29. The composition of Claim 26 wherein the platelet aggregation inhibitor is selected from the group consisting of ticlopidine and clopidogrel.

30. The composition of Claim 26 wherein the platelet aggregation inhibitor is a glycoprotein IIb/IIIa receptor antagonist.

25 31. The composition of Claim 30 wherein the glycoprotein IIb/IIIa receptor antagonist is selected from the group consisting of:

30 Mpr-(Acetimidyl-Lys)-Gly-Asp-Trp-Phe-Cys-NH₂,
Mpr-(Acetimidyl-Lys)-Gly-Asp-Trp-Phe-Pen-NH₂,
Mpr-(Phenylimidyl-Lys)-Gly-Asp-Trp-Phe-Pen-NH₂,
Mpr-(Phenylimidyl-Lys)-Gly-Asp-Trp-Phe-Cys-NH₂,
N-Methyl-D-phenylalanyl-N-[(1S)-1-formyl-4-guanidinobutyl]-L-prolinamide,

- 34 -

((1-(2-((4-(aminoiminomethyl)benzoyl)amino)-3-(4-hydroxyphenyl)-1-oxopropyl)-4-piperidinyl)oxy)-(S)-acetic acid,
 N-(2-(2-(((3-((aminoiminomethyl)amino)propyl)amino)carbonyl)-1-piperidnyl)-1-(cyclohexylmethyl)-2-oxoethyl)-(R,S)-glycine,
 5 Ethyl 3-[[4-[[4-(aminoiminomethyl)phenyl]amino]-1,4-dioxobutyl]amino]-4-pentynoate,
 (2-S-(n-Butylsulfonylamino)-3[4-(piperidin-4-yl)butyloxyphenyl]propionic acid hydrochloride, and
 2(S)-[(p-Toluenesulfonyl)amino]-3-[[[5,6,7,8-tetrahydro-4-oxo-5-[2-(piperidin-4-yl)ethyl]-4H-pyrazolo-[1,5-a][1,4]diazepin-2-yl]carbonyl]-amino]propionic acid;
 10 and the pharmaceutically acceptable salts, esters and pro-drugs thereof.

32. The composition of Claim 30 wherein the a
 15 glycoprotein IIb/IIIa receptor antagonist is selected from the group consisting of: DMP 754, [3(R)-[2-piperidin-4-yl)ethyl]-2-piperidone-1]acetyl-3(R)-methyl- β -alanine, and 5-[(4-Piperidinyl)methoxy]-2-indolecarbonyl-2(S)-phenylsulfonyl-amino- β -alanine and the pharmaceutically acceptable salts, esters and pro-drugs thereof.

20

33. The composition of Claim 32 wherein the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin and the pharmaceutically acceptable salts and esters thereof.

25

34. The composition of Claim 26 adapted for oral administration.

35. A pharmaceutical composition made by combining a platelet aggregation inhibitor, an HMG-CoA reductase inhibitor and a
 30 pharmaceutically acceptable carrier.

36. The composition of Claim 35 wherein the platelet aggregation inhibitor is selected from the group consisting of a

- 35 -

glycoprotein IIb/IIIa receptor antagonist, ticlopidine, clopidogrel, aspirin and dipyridamole.

37. A process for making a pharmaceutical composition
5 comprising combining a platelet aggregation inhibitor, an HMG-CoA reductase inhibitor and a pharmaceutically acceptable carrier.

38. The process of Claim 37 wherein the platelet
10 aggregation inhibitor is selected from the group consisting of a glycoprotein IIb/IIIa receptor antagonist, ticlopidine, clopidogrel, aspirin and dipyridamole.

39. The method of Claim 2 wherein the glycoprotein
IIb/IIIa receptor antagonist is selected from the group consisting of:
15 [3(R)-[2-piperidin-4-yl)ethyl]-2-piperidone-1]acetyl-3(R)-methyl- β -alanine;
5-[(4-Piperidinyl)methoxy]-2-indolecarbonyl-2(S)-phenylsulfonyl-amino- β -alanine;
2(S)-[(p-Toluenesulfonyl)amino]-3-[[[5,6,7,8-tetrahydro-4-oxo-5-[2-
20 (piperidin-4-yl)ethyl]-4H-pyrazolo-[1,5-a][1,4]diazepin-2-yl]carbonyl]-amino]propionic acid;
MK-383; DMP 728; DMP 754; Ro44-9883; Ro43-8857; sibrafiban; xemlofiban; fradafiban; SB 214857; ZD-2486; TAK-029; orbofiban; RPR 109891; GR144053; GR233548; and SDZ 562; and the
25 pharmaceutically acceptable salts, esters and pro-drugs thereof.

40. The composition of Claim 30 wherein the
glycoprotein IIb/IIIa receptor antagonist is selected from the group
consisting of :
30 [3(R)-[2-piperidin-4-yl)ethyl]-2-piperidone-1]acetyl-3(R)-methyl- β -alanine;
5-[(4-Piperidinyl)methoxy]-2-indolecarbonyl-2(S)-phenylsulfonyl-amino- β -alanine;

- 36 -

2(S)-[(p-Toluenesulfonyl)amino]-3-[[[5,6,7,8-tetrahydro-4-oxo-5-[2-(piperidin-4-yl)ethyl]-4H-pyrazolo-[1,5-a][1,4]diazepin-2-yl]carbonyl]-amino]propionic acid;

5 MK-383; DMP 728; DMP 754; Ro44-9883; Ro43-8857; sibrafiban; xemlofiban; fradafiban; SB 214857; ZD-2486; TAK-029; orbofiban; RPR 109891; GR144053; GR233548; and SDZ 562; and the pharmaceutically acceptable salts, esters and pro-drugs thereof.

41. A kit comprised of an HMG-CoA reductase inhibitor
10 in an oral dosage formulation and a platelet aggregation inhibitor in a separate oral dosage formulation.

42. The kit of claim 41 wherein the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin, and the platelet
15 aggregation inhibitor is a GP IIb/IIIa receptor antagonist.

43. The kit of claim 41 wherein the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin, and the platelet aggregation inhibitor is aspirin.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/16389

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/505, 31/44, 31/35, 31/225

US CL : 514/258, 301, 460, 548

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/258, 301, 460, 548

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS AND CAS ONLINE: ischemi? or thrombo? or resteno? or infarc? with hmg-coa reductase inhibitor or platelet aggregation inhibitor

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,140,012 A (MCGOVERN et al.) 18 August 1992, see abstract.	1-43
Y	Chemical Abstract No. 123:13273, 1995, HIRSCH. 'HMG-CoA Reductase Inhibitors in the Normalization of Vascular Endothelial Dysfunction.	1-43
Y	Chemical Abstract No. 118:94100, 1993, ROUX. 'Effects of Heparin, Aspirin and a Synthetic Platelet Glycoprotein IIb-IIIa Receptor Antagonist (Ro 43-5054) on Coronary Artery Reperfusion and Reocclusion After Thrombolysis with Tissue-Type Plasminogen Activator in the Dog.	1-43

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 06 NOVEMBER 1997	Date of mailing of the international search report 06 JAN 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Kimberly Jordan</i> KIMBERLY JORDAN Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/16388

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Chemical Abstract No. 101:83760, 1984, FAXON. 'Effect of Antiplatelet Therapy on Restenosis After Experimental Angioplasty.'	1-43